



Contents lists available at ScienceDirect

Environmental Research

journal homepage: www.elsevier.com/locate/envres

5 G wireless telecommunications expansion: Public health and environmental implications[☆]

Cindy L. Russell

PO Box 7443, Menlo Park, CA 94026, USA

ARTICLE INFO

Keywords:

Wireless technology
Public health
Radiofrequency radiation
Environmental pollution

ABSTRACT

The popularity, widespread use and increasing dependency on wireless technologies has spawned a telecommunications industrial revolution with increasing public exposure to broader and higher frequencies of the electromagnetic spectrum to transmit data through a variety of devices and infrastructure. On the horizon, a new generation of even shorter high frequency 5G wavelengths is being proposed to power the Internet of Things (IoT). The IoT promises us convenient and easy lifestyles with a massive 5G interconnected telecommunications network, however, the expansion of broadband with shorter wavelength radiofrequency radiation highlights the concern that health and safety issues remain unknown. Controversy continues with regards to harm from current 2G, 3G and 4G wireless technologies. 5G technologies are far less studied for human or environmental effects.

It is argued that the addition of this added high frequency 5G radiation to an already complex mix of lower frequencies, will contribute to a negative public health outcome both from both physical and mental health perspectives.

Radiofrequency radiation (RF) is increasingly being recognized as a new form of environmental pollution. Like other common toxic exposures, the effects of radiofrequency electromagnetic radiation (RF EMR) will be problematic if not impossible to sort out epidemiologically as there no longer remains an unexposed control group. This is especially important considering these effects are likely magnified by synergistic toxic exposures and other common health risk behaviors. Effects can also be non-linear. Because this is the first generation to have cradle-to-grave lifespan exposure to this level of man-made microwave (RF EMR) radiofrequencies, it will be years or decades before the true health consequences are known. Precaution in the roll out of this new technology is strongly indicated.

This article will review relevant electromagnetic frequencies, exposure standards and current scientific literature on the health implications of 2G, 3G, 4G exposure, including some of the available literature on 5G frequencies. The question of what constitutes a public health issue will be raised, as well as the need for a precautionary approach in advancing new wireless technologies.

1. Introduction

The adoption of new 5G technology promises to give the public a transformative communication network with an explosion of speed, volume of data and number of devices with unlimited computing instantly to anyone in the world. High tech companies are already marketing the Internet of Things to businesses, healthcare systems, schools and the public. The promise to connect our phones and appliances, will virtually eliminate many day-to-day household and business functions including driving. This will, according to industry, create a superior, connected society and unprecedented economic growth. What is missing in this discussion is the maturing literature on adverse

biological, physiological, and psychological health effects of the 2G, 3G, and 4G radiofrequencies we are already exposed to, in addition to indications from the scientific literature that 5G frequencies could also be hazardous.

Many important but unanswered questions merit serious consideration. Is the widespread deployment of this pervasive higher frequency small cell distributed antennae system in our cities and on our homes safe for humans and the environment? Will it add to the burden of chronic disease that costs our nation, according to the CDC, an estimated 2.3 trillion dollars annually (CDC, 2017)? Are we already digitally over connected, shrinking our gray matter and becoming a dysfunctional addicted nation because of it (Weng et al., 2012)? How

[☆] The author has read and approved the paper and it has not been published previously nor is it being considered by any other peer-reviewed journal. The author has no conflicts of interest. There was no financial support for the research, design, analysis, interpretation or writing of this article nor for the decision to publish this article.

E-mail address: cindyleerussell@gmail.com.

<https://doi.org/10.1016/j.envres.2018.01.016>

Received 3 October 2017; Accepted 16 January 2018

0013-9351/ © 2018 Elsevier Inc. All rights reserved.



Contents lists available at ScienceDirect

Environmental Research

journal homepage: www.elsevier.com/locate/envres

Report of final results regarding brain and heart tumors in Sprague-Dawley rats exposed from prenatal life until natural death to mobile phone radiofrequency field representative of a 1.8 GHz GSM base station environmental emission

L. Falcioni, L. Bua, E. Tibaldi, M. Lauriola, L. De Angelis, F. Gnudi, D. Mandrioli, M. Manservigi, F. Manservigi, I. Manzoli, I. Menghetti, R. Montella, S. Panzacchi, D. Sgargi, V. Strollo, A. Vornoli, F. Belpoggi*

Cesare Maltoni Cancer Research Center, Ramazzini Institute, Castello di Bentivoglio, via Saliceto 3, Bentivoglio, 40010 Bologna, Italy

ARTICLE INFO

Keywords:

RF
Sprague-Dawley rats
Life-span bioassay
Mobile phone
Carcinogenicity

ABSTRACT

Background: In 2011, IARC classified radiofrequency radiation (RFR) as possible human carcinogen (Group 2B). According to IARC, animals studies, as well as epidemiological ones, showed limited evidence of carcinogenicity. In 2016, the NTP published the first results of its long-term bioassays on near field RFR, reporting increased incidence of malignant glial tumors of the brain and heart Schwannoma in rats exposed to GSM – and CDMA – modulated cell phone RFR. The tumors observed in the NTP study are of the type similar to the ones observed in some epidemiological studies of cell phone users.

Objectives: The Ramazzini Institute (RI) performed a life-span carcinogenic study on Sprague-Dawley rats to evaluate the carcinogenic effects of RFR in the situation of far field, reproducing the environmental exposure to RFR generated by 1.8 GHz GSM antenna of the radio base stations of mobile phone. This is the largest long-term study ever performed in rats on the health effects of RFR, including 2448 animals. In this article, we reported the final results regarding brain and heart tumors.

Methods: Male and female Sprague-Dawley rats were exposed from prenatal life until natural death to a 1.8 GHz GSM far field of 0, 5, 25, 50 V/m with a whole-body exposure for 19 h/day.

Results: A statistically significant increase in the incidence of heart Schwannomas was observed in treated male rats at the highest dose (50 V/m). Furthermore, an increase in the incidence of heart Schwann cells hyperplasia was observed in treated male and female rats at the highest dose (50 V/m), although this was not statistically significant. An increase in the incidence of malignant glial tumors was observed in treated female rats at the highest dose (50 V/m), although not statistically significant.

Conclusions: The RI findings on far field exposure to RFR are consistent with and reinforce the results of the NTP study on near field exposure, as both reported an increase in the incidence of tumors of the brain and heart in RFR-exposed Sprague-Dawley rats. These tumors are of the same histotype of those observed in some epidemiological studies on cell phone users. These experimental studies provide sufficient evidence to call for the re-evaluation of IARC conclusions regarding the carcinogenic potential of RFR in humans.

1. Introduction

Early warnings on the potential carcinogenic risks of mobile phone radiofrequency radiation (RFR) raised in the early 2000 when, for the first time, it was published that people using mobile phones had a significant increased risk to develop vestibular Schwannoma and brain tumors (Hardell et al., 2003, 2002). In 2011, the International Agency

for Research on Cancer (IARC) classified RFR as possible human carcinogen (Group 2B) based on limited evidence both in humans and experimental animals (Baan et al., 2011; IARC, 2013). Two epidemiological case-control studies resulted more informative for the IARC evaluation, showing that the risk to develop brain tumors and vestibular Schwannoma was increased in people with the highest cumulative use of mobile phones, in people who had used mobile phones on the

* Corresponding author.

E-mail address: belpoggif@ramazzini.it (F. Belpoggi).

<https://doi.org/10.1016/j.envres.2018.01.037>

Received 9 October 2017; Received in revised form 19 December 2017; Accepted 23 January 2018
0013-9351/ © 2018 Elsevier Inc. All rights reserved.