


Contents lists available at ScienceDirect

Toxicology Letters





Toxicology and cellular mechanisms of electromagnetic fields (EMF)-Health aspects of exposure to EMF Emitted by wireless mobile systems and emerging technologies

S12-01 Two-year oncogenicity evaluations of cell phone radiofrequency radiation in Sprague-Dawley rats and B6C3F1 mice



David McCormick

IIT Research Institute, Chicago, IL, United States

Epidemiology data concerning possible health effects of exposure to radiofrequency fields (RF) are conflicting. For this reason, welldesigned and controlled studies in predictive laboratory animal models provide the best prospective opportunity to identify effects of RF exposure that may translate into human health hazards. The U.S. National Toxicology Program supported a program in our laboratory to identify and characterize effects of acute, subchronic, and chronic exposure to non-thermal levels of RF in Sprague-Dawley rats and B6C3F1 mice. Five-day pilot studies were performed to identify the maximum Specific Absorption Ratios (SARs) to which juvenile, adult, and pregnant rodents can be exposed without increasing body temperature by >1.0 °C. Subsequent subchronic (ten-week) toxicity studies failed to identify any toxicologically significant effects of non-thermal RF on survival, body weight, clinical signs, hematology, or gross or microscopic pathology. Two-year studies were performed to determine if exposure to non-thermal levels of RF increases the incidence of neoplasia in any site. Male rats exposed to RF demonstrated significantly increased incidences of glioma (brain) and schwannoma (heart); these increases were not seen in female rats or in either sex of mice. Gliomas and schwannomas have been identified in some epidemiology studies as possible RF-induced neoplasms. Considering (a) the conflicting results of RF epidemiology studies and (b) the lack of generally accepted biophysical or molecular mechanisms through which RF could induce or promote neoplasia, data from animal bioassays will play a central role in "weight-of-theevidence" assessments of the possible health effects of RF exposure.

Supported by HHSN29120055544.

http://dx.doi.org/10.1016/j.toxlet.2017.07.075